







Polar-koordinater

<p>Koordinater</p>	$x = r \cos \theta$ $y = r \sin \theta$ $x^2 + y^2 = r^2$	
<p>Areal</p>	$A = \int_{\theta_1}^{\theta_2} \frac{1}{2} r^2 d\theta$	 
<p>Lengden av polar kurve</p>	$L = \int_{\theta_1}^{\theta_2} \sqrt{r^2 + \left(\frac{dr}{d\theta}\right)^2} d\theta$	<p> $e < 1$ Ellipse $e = 1$ Parabel $e > 1$ Hyperbel </p> 
<p>Polar ligning for konisk seksjon</p>	$r = \frac{ke}{1 + e \cos \theta}$	<p>Ellipse</p>



Symmetri om x-aksen:

$$r(-\theta) = r(\theta) \quad \vee \quad r(\pi - \theta) = -r(\theta)$$





Symmetri om y-aksen:

$$r(-\theta) = -r(\theta) \quad \vee \quad r(\pi - \theta) = r(\theta)$$







Symmetri om origo:






$$r(\theta) = -r(\theta) \quad \vee \quad r(\pi + \theta) = r(\theta)$$







Vektorer og geometri i rommet - Vektor

<p>Vektor</p>	$\vec{v} = (x_2 - x_1)\vec{i} + (y_2 - y_1)\vec{j} + (z_2 - z_1)\vec{k}$ $= [x_2 - x_1, y_2 - y_1, z_2 - z_1]$ $= [v_1, v_2, v_3]$	
<p>Lengden av en vektor</p>	$ \vec{v} = \sqrt{v_1^2 + v_2^2 + v_3^2}$	
<p>Enhetsvektor</p>	$\vec{n} = \frac{\vec{v}}{ \vec{v} }$	
<p>Addisjon</p>	$\vec{u} = [u_1, u_2, u_3]$ $\vec{v} = [v_1, v_2, v_3]$ $\vec{u} + \vec{v} = [u_1 + v_1, u_2 + v_2, u_3 + v_3]$	
<p>Multiplikasjon med skalar</p>	$\vec{u} = [u_1, u_2, u_3]$ $k\vec{u} = [ku_1, ku_2, ku_3]$	
<p>Regneregler for vektorer</p>	$\vec{u} + \vec{v} = \vec{v} + \vec{u}$ $(\vec{u} + \vec{v}) + \vec{w} = \vec{u} + (\vec{v} + \vec{w})$ $\vec{u} + \vec{0} = \vec{u}$ $\vec{u} + (-\vec{u}) = \vec{0}$ $0\vec{u} = \vec{0}$ $1\vec{u} = \vec{u}$ $a(b\vec{u}) = (ab)\vec{u}$ $a(\vec{u} + \vec{v}) = a\vec{u} + a\vec{v}$ $(a + b)\vec{u} = a\vec{u} + b\vec{u}$	





Vektorer og geometri i rommet - Skalarprodukt

Skalarprodukt	$\vec{u} \cdot \vec{v} = \vec{u} \vec{v} \cos \theta = u_1v_1 + u_2v_2 + u_3v_3$	
Regler	$\vec{u} \cdot \vec{v} = \vec{v} \cdot \vec{u}$ $(c\vec{u}) \cdot \vec{v} = \vec{u} \cdot (c\vec{v})$ $\vec{u} \cdot (\vec{v} + \vec{w}) = \vec{u} \cdot \vec{v} + \vec{u} \cdot \vec{w}$ $\vec{u} \cdot \vec{u} = \vec{u} ^2$ $\vec{0} \cdot \vec{u} = 0$	
Ortogonale vektorer	$\vec{u} \perp \vec{v} \Leftrightarrow \vec{u} \cdot \vec{v} = 0$	
Projeksjon av u-vektor på v-vektor Skalarkomponent av u-vektor i retning v-vektor	$proj_{\vec{v}} \vec{u} = \frac{\vec{u} \cdot \vec{v}}{ \vec{v} ^2} \vec{v}$ $ \vec{u} \cos \theta = \vec{u} \cdot \frac{\vec{v}}{ \vec{v} }$	
Vektor skrevet som en sum av ortogonale vektorer	$\vec{u} = \underbrace{\frac{\vec{u} \cdot \vec{v}}{ \vec{v} ^2} \vec{v}}_{\text{Parallell med } \vec{v}} + \underbrace{\left(\vec{u} - \frac{\vec{u} \cdot \vec{v}}{ \vec{v} ^2} \vec{v} \right)}_{\text{Normal på } \vec{v}}$	




Vektorer og geometri i rommet - Vektorprodukt

<p>Vektorprodukt</p>	$\vec{u} \times \vec{v} = \ \vec{u}\ \ \vec{v}\ \sin \theta \vec{n}$ $= \begin{vmatrix} \vec{i} & \vec{j} & \vec{k} \\ u_1 & u_2 & u_3 \\ v_1 & v_2 & v_3 \end{vmatrix}$ $= [u_2v_3 - u_3v_2, u_3v_1 - u_1v_3, u_1v_2 - u_2v_1]$	
<p>Regler</p>	$(r\vec{u}) \times (s\vec{v}) = (rs)\vec{u} \times \vec{v}$ $\vec{u} \times (\vec{v} + \vec{w}) = \vec{u} \times \vec{v} + \vec{u} \times \vec{w}$ $(\vec{v} + \vec{w}) \times \vec{u} = \vec{v} \times \vec{u} + \vec{w} \times \vec{u}$ $\vec{v} \times \vec{u} = -(\vec{u} \times \vec{v})$ $\vec{0} \times \vec{u} = \vec{0}$	
	$\vec{i} \times \vec{j} = \vec{k}$ $\vec{j} \times \vec{k} = \vec{i}$ $\vec{k} \times \vec{i} = \vec{j}$	
<p>Parallele vektorer</p>	$\vec{u} \parallel \vec{v} \Leftrightarrow \vec{u} \times \vec{v} = \vec{0}$	





Vektorer og geometri i rommet - Areal - Volum

<p>Areal</p>	$A = \vec{u} \times \vec{v} $	
<p>Volum</p>	$V = (\vec{u} \times \vec{v}) \cdot \vec{w} $ $= \begin{vmatrix} u_1 & u_2 & u_3 \\ v_1 & v_2 & v_3 \\ w_1 & w_2 & w_3 \end{vmatrix}$	
	$\vec{i} \times \vec{j} = \vec{k}$ $\vec{j} \times \vec{k} = \vec{i}$ $\vec{k} \times \vec{i} = \vec{j}$	
<p>Parallele vektorer</p>	$\vec{u} \parallel \vec{v} \Leftrightarrow \vec{u} \times \vec{v} = \vec{0}$	




Vektorer og geometri i rommet - Linjer - Plan

<p>Linje gjennom P_0 parallell med v-vektor</p>	$\vec{r}(t) = \vec{r}_0 + t\vec{v}$ $[x, y, z] = [x_0, y_0, z_0] + t[v_1, v_2, v_3]$ $x = x_0 + tv_1$ $y = y_0 + tv_2$ $z = z_0 + tv_3$	
<p>Avstand fra et punkt S til en linje gjennom P parallell med v-vektor</p>	$d = \frac{ \vec{PS} \times \vec{v} }{ \vec{v} }$	
<p>Plan gjennom $P_0(x_0, y_0, z_0)$ med normalvektor $\vec{n} = [A, B, C]$</p>	$A(x - x_0) + B(y - y_0) + C(z - z_0) = 0$ $Ax + By + Cz = D \quad D = Ax_0 + By_0 + Cz_0$	

Vektorer og geometri i rommet - Kvadratiske flater

<p>Kvadratiske flater</p>	$Ax^2 + By^2 + Cz^2 + Dxy + Eyz + Fxz + Gx + Hy + Jz + K = 0$	
<p>Parabolisk sylinder</p>	$y = x^2$	
<p>Elliptisk sylinder</p>	$\frac{x^2}{a^2} + \frac{z^2}{b^2} = 1$	
<p>Ellipsoide</p>	$\frac{x^2}{a^2} + \frac{y^2}{b^2} + \frac{z^2}{c^2} = 1$	
<p>Elliptisk paraboloid</p>	$\frac{x^2}{a^2} + \frac{y^2}{b^2} = \frac{z}{c}$	

Vektorer og geometri i rommet - Kvadratiske flater


<p>Elliptisk kjegle</p>	$\frac{x^2}{a^2} + \frac{y^2}{b^2} = \frac{z^2}{c^2}$	
<p>Hyperboloide</p>	$\frac{x^2}{a^2} + \frac{y^2}{b^2} - \frac{z^2}{c^2} = 1$ <p style="text-align: right;">En flate</p>	
<p>Hyperboloide</p>	$\frac{z^2}{c^2} - \frac{x^2}{a^2} - \frac{y^2}{b^2} = 1$ <p style="text-align: right;">To flater</p>	

Vektor-funksjoner

Posisjon	$\vec{r}(t)$	
Hastighet	$\vec{v}(t) = \vec{r}'(t) = \frac{d\vec{r}(t)}{dt}$	Fart $\frac{ds}{dt} = v(t) = \vec{v}(t) $
Akselerasjon	$\vec{a}(t) = \vec{v}'(t) = \vec{r}''(t) = \frac{d^2\vec{r}(t)}{dt^2}$	
Enhets tangent vektor	$\vec{T} = \frac{d\vec{r}}{ds} = \frac{\vec{v}}{ \vec{v} }$	
Krumning	$\kappa = \left \frac{d\vec{T}}{ds} \right = \frac{1}{ \vec{v} } \left \frac{d\vec{T}}{dt} \right = \frac{ \vec{v} \times \vec{a} }{ \vec{v} ^3}$	
Hoved enhets normal	$\vec{N} = \frac{1}{\kappa} \frac{d\vec{T}}{ds} = \frac{\frac{d\vec{T}}{dt}}{\left \frac{d\vec{T}}{dt} \right }$	
Binormal	$\vec{B} = \vec{T} \times \vec{N}$	
Torsjon	$\tau = -\frac{d\vec{B}}{ds} \cdot \vec{N} = \frac{\begin{vmatrix} \dot{x} & \dot{y} & \dot{z} \\ \ddot{x} & \ddot{y} & \ddot{z} \end{vmatrix}}{ \vec{v} \times \vec{a} ^2}$	
Akselerasjonskomponenter	$\vec{a} = a_T \vec{T} + a_N \vec{N}$	$a_T = \frac{d}{dt} \vec{v} $ $a_N = \kappa \vec{v} ^2 = \sqrt{ \vec{a} ^2 - a_T^2}$

Koordinatsystemer

<p>Sylindrisk → Rektangulær</p>	$x = r \cos \theta$ $y = r \sin \theta$ $z = z$
<p>Sfærisk → Rektangulær</p>	$x = \rho \sin \varphi \cos \theta$ $y = \rho \sin \varphi \sin \theta$ $z = \rho \cos \varphi$
<p>Sfærisk → Sylindrisk</p>	$r = \rho \sin \varphi$ $\theta = \theta$ $z = \rho \cos \varphi$
<p>Rektangulær</p> <p>Sylindrisk</p> <p>Sfærisk</p>	$dV = dx dy dz$ $dV = dz r dr d\theta$ $dV = \rho^2 \sin \varphi d\rho d\varphi d\theta$



Vektor-felt - Del-operator

Funksjon	$f = f(x, y, z)$
Vektor-felt	$\vec{F} = [F_1, F_2, F_3]$
Potensial-funksjon f	$\vec{F} = \nabla f$
Del-operator	$\nabla = \left[\frac{\partial}{\partial x}, \frac{\partial}{\partial y}, \frac{\partial}{\partial z} \right]$
Gradient	$\nabla f = \left[\frac{\partial f}{\partial x}, \frac{\partial f}{\partial y}, \frac{\partial f}{\partial z} \right]$
Divergens	$\text{div} \vec{F} = \nabla \cdot \vec{F} = \left[\frac{\partial}{\partial x}, \frac{\partial}{\partial y}, \frac{\partial}{\partial z} \right] \cdot [F_1, F_2, F_3] = \frac{\partial F_1}{\partial x} + \frac{\partial F_2}{\partial y} + \frac{\partial F_3}{\partial z}$
Curl	$\text{curl} \vec{F} = \nabla \times \vec{F} = \left[\frac{\partial}{\partial x}, \frac{\partial}{\partial y}, \frac{\partial}{\partial z} \right] \times [M, N, P] = \begin{vmatrix} \vec{i} & \vec{j} & \vec{k} \\ \frac{\partial}{\partial x} & \frac{\partial}{\partial y} & \frac{\partial}{\partial z} \\ F_1 & F_2 & F_3 \end{vmatrix} = \left[\frac{\partial F_3}{\partial y} - \frac{\partial F_2}{\partial z}, \frac{\partial F_1}{\partial z} - \frac{\partial F_3}{\partial x}, \frac{\partial F_2}{\partial x} - \frac{\partial F_1}{\partial y} \right]$

Kurve-integral

Kurve-integral	$\int_C f(x, y, z) ds = \int_a^b f(g(t), h(t), k(t)) \vec{r}'(t) dt$	Kurve-lengde $\int_C ds = \int_a^b \vec{r}'(t) dt$										
Arbeid	$W = \int_C \vec{F} \cdot \vec{T} ds = \int_C \vec{F} \cdot d\vec{r} = \int_C F_1 dx + F_2 dy + F_3 dz$											
Strømning Sirkulasjon	$\int_C \vec{F} \cdot \vec{T} ds = \int_C \vec{F} \cdot d\vec{r} = \int_C F_1 dx + F_2 dy$ $\oint_C \vec{F} \cdot \vec{T} ds = \oint_C \vec{F} \cdot d\vec{r} = \oint_C F_1 dx + F_2 dy$	I planet										
Fluks	$\int_C \vec{F} \cdot \vec{n} ds = \int_C F_1 dy - F_2 dx \quad \vec{n} = \vec{T} \times \vec{k}$	I planet										
Fundamental-teoremet for kurve-integraler	\vec{F} vektorfelt med kontinuerlige komponenter over et åpent område D \Downarrow \exists differensierbar funksjon $f: \vec{F} = \nabla f$ $\int_A^B \vec{F} \cdot d\vec{r} = f(B) - f(A)$											
Eksakt differensialform	$F_1 dx + F_2 dy + F_3 dz$ er en differensialform En differensialform er eksakt hvis $M dx + N dy + P dz = \frac{\partial f}{\partial x} dx + \frac{\partial f}{\partial y} dy + \frac{\partial f}{\partial z} dz$ Differensialform $F_1 dx + F_2 dy + F_3 dz$ er eksakt \Updownarrow $\frac{\partial F_3}{\partial y} = \frac{\partial F_2}{\partial z} \quad \frac{\partial F_1}{\partial z} = \frac{\partial F_3}{\partial x} \quad \frac{\partial F_2}{\partial x} = \frac{\partial F_1}{\partial y}$											
F konservativ (vei-uavhengig)	$\int_C \vec{F} \cdot d\vec{r} = f(b) - f(a)$ $\oint_C \vec{F} \cdot d\vec{r} = 0$ $\vec{F} = \nabla f$ $\text{curl} \vec{F} = \nabla \times \vec{F} = \vec{0}$ <table style="width: 100%; border: none;"> <tr> <td style="text-align: center;">$\frac{\partial F_3}{\partial y} = \frac{\partial F_2}{\partial z}$</td> <td style="text-align: center;">$\frac{\partial F_1}{\partial z} = \frac{\partial F_3}{\partial x}$</td> <td style="text-align: center;">$\frac{\partial F_2}{\partial x} = \frac{\partial F_1}{\partial y}$</td> <td style="text-align: center;">\vec{F}</td> <td style="text-align: center;">Konservativ</td> </tr> <tr> <td style="text-align: center;">$\frac{\partial F_3}{\partial y} = \frac{\partial F_2}{\partial z}$</td> <td style="text-align: center;">$\frac{\partial F_1}{\partial z} = \frac{\partial F_3}{\partial x}$</td> <td style="text-align: center;">$\frac{\partial F_2}{\partial x} = \frac{\partial F_1}{\partial y}$</td> <td style="text-align: center;">$F_1 dx + F_2 dy + F_3 dz$</td> <td style="text-align: center;">Eksakt</td> </tr> </table>		$\frac{\partial F_3}{\partial y} = \frac{\partial F_2}{\partial z}$	$\frac{\partial F_1}{\partial z} = \frac{\partial F_3}{\partial x}$	$\frac{\partial F_2}{\partial x} = \frac{\partial F_1}{\partial y}$	\vec{F}	Konservativ	$\frac{\partial F_3}{\partial y} = \frac{\partial F_2}{\partial z}$	$\frac{\partial F_1}{\partial z} = \frac{\partial F_3}{\partial x}$	$\frac{\partial F_2}{\partial x} = \frac{\partial F_1}{\partial y}$	$F_1 dx + F_2 dy + F_3 dz$	Eksakt
$\frac{\partial F_3}{\partial y} = \frac{\partial F_2}{\partial z}$	$\frac{\partial F_1}{\partial z} = \frac{\partial F_3}{\partial x}$	$\frac{\partial F_2}{\partial x} = \frac{\partial F_1}{\partial y}$	\vec{F}	Konservativ								
$\frac{\partial F_3}{\partial y} = \frac{\partial F_2}{\partial z}$	$\frac{\partial F_1}{\partial z} = \frac{\partial F_3}{\partial x}$	$\frac{\partial F_2}{\partial x} = \frac{\partial F_1}{\partial y}$	$F_1 dx + F_2 dy + F_3 dz$	Eksakt								

Flate-integral

Flate-integral	$\iint_S g dS = \iint_R g(x, y, z) \frac{ \nabla f }{ \nabla f \cdot \vec{p} } dA$	Flate-areal $\iint_S dS = \iint_R \frac{ \nabla f }{ \nabla f \cdot \vec{p} } dA$
Parameterisert flate-integral	$\iint_S G(x, y, z) dS = \iint_R G(f(u, v), g(u, v), h(u, v)) \vec{r}_u \times \vec{r}_v dudv$	Flate-areal $\iint_S dS = \iint_R \vec{r}_u \times \vec{r}_v dudv$
Fluks	$\iint_S \vec{F} \cdot \vec{n} dS$	
Enhetsnormalvektor	$\vec{n} = \frac{\nabla f}{ \nabla f }$ f skalarfunksjon som har flaten S som en nivåflate	$\vec{n} = \frac{\vec{r}_u \times \vec{r}_v}{ \vec{r}_u \times \vec{r}_v }$ Parameterisert flate

Green's teorem - Stoke's teorem


	2D	3D	
Green - Normalform Divergens = Flukstetthet $= \frac{\text{Fluks}}{\text{Areal}}$	$\oint_C \vec{F} \cdot \vec{n} ds = \oint_C F_1 dy - F_2 dx$ $= \iint_R \left[\frac{\partial F_1}{\partial x} + \frac{\partial F_2}{\partial y} \right] dx dy$ $= \iint_R \text{div} \vec{F} dA$ $= \iint_R \nabla \cdot \vec{F} dA$	$\iiint_S \vec{F} \cdot \vec{n} dS = \iiint_S F_1 dy dz + F_2 dz dx + F_3 dx dy$ $= \iiint_D \left[\frac{\partial F_1}{\partial x} + \frac{\partial F_2}{\partial y} + \frac{\partial F_3}{\partial z} \right] dx dy dz$ $= \iiint_D \text{div} \vec{F} dV$ $= \iiint_D \nabla \cdot \vec{F} dV$	Gauss Divergens
Green - Tangensialform Curl = Sirkulasjonstetthet $= \frac{\text{Sirkulasjon}}{\text{Areal}}$	$\oint_C \vec{F} \cdot \vec{T} ds = \oint_C \vec{F} \cdot d\vec{r}$ $= \oint_C F_1 dx + F_2 dy$ $= \iint_R \left[\frac{\partial F_2}{\partial x} - \frac{\partial F_1}{\partial y} \right] dx dy$ $= \iint_R \text{curl} \vec{F} \cdot \vec{k} dA$ $= \iint_R (\nabla \times \vec{F}) \cdot \vec{k} dA$	$\oint_C \vec{F} \cdot \vec{T} ds = \oint_C \vec{F} \cdot d\vec{r}$ $= \oint_C F_1 dx + F_2 dy + F_3 dz$ $= \iint_S \left(\frac{\partial F_3}{\partial y} - \frac{\partial F_2}{\partial z} \right) dy dz + \left(\frac{\partial F_1}{\partial z} - \frac{\partial F_3}{\partial x} \right) dz dx + \left(\frac{\partial F_2}{\partial x} - \frac{\partial F_1}{\partial y} \right) dx dy$ $= \iint_S \text{curl} \vec{F} \cdot \vec{n} dS$ $= \iint_S \nabla \times \vec{F} \cdot \vec{n} d\sigma$	Stoke

2D**Green - Divergens**

$$\Phi = \oint_C \vec{F} \cdot \vec{n} ds = \iint_R \nabla \cdot \vec{F} dA$$

Green - Curl


$$C = \oint_C \vec{F} \cdot \vec{T} ds = \iint_R (\nabla \times \vec{F}) \cdot \vec{k} dA$$

3D**Gauss - Divergens**

$$\Phi = \oint_S \vec{F} \cdot \vec{n} dS = \iiint_D \nabla \cdot \vec{F} dV$$

Stokes - Curl

$$C = \oint_C \vec{F} \cdot \vec{T} ds = \iint_S (\nabla \times \vec{F}) \cdot \vec{n} dS$$



Substitusjon i multiple integraler

<p>2D</p>	$\iint_D F(x, y) dx dy = \iint_D F(x(u, v), y(u, v)) J(u, v) du dv$ $J(u, v) = \begin{vmatrix} x_u & x_v \\ y_u & y_v \end{vmatrix} \quad J^{-1}(x, y) = \begin{vmatrix} u_x & u_y \\ v_x & v_y \end{vmatrix} \quad J(u, v) J^{-1}(x, y) = 1$
<p>3D</p>	$\iiint_D F(x, y, z) dx dy dz = \iiint_G F(x(u, v, w), y(u, v, w), z(u, v, w)) J(u, v, w) du dv dw$ $J(u, v, w) = \begin{vmatrix} x_u & x_v & x_w \\ y_u & y_v & y_w \\ z_u & z_v & z_w \end{vmatrix} \quad J^{-1}(x, y, z) = \begin{vmatrix} u_x & u_y & u_z \\ v_x & v_y & v_z \\ w_x & w_y & w_z \end{vmatrix} \quad J(u, v, w) J^{-1}(x, y, z) = 1$

Greens form av areal

<p>Tangensiell form</p>	$A = \iint dA = \oint x dy \quad \oint F_1 dx + F_2 dy = \iint \left[\frac{\partial F_2}{\partial x} - \frac{\partial F_1}{\partial y} \right] dA \quad F_2 = x \quad F_1 = 0$ $A = \iint dA = -\oint y dx \quad \oint F_1 dx + F_2 dy = \iint \left[\frac{\partial F_2}{\partial x} - \frac{\partial F_1}{\partial y} \right] dA \quad F_2 = 0 \quad F_1 = -y$
<p>Normalform</p>	$A = \iint dA = \frac{1}{2} \oint x dy - y dx \quad \oint F_1 dy - F_2 dx = \iint \left[\frac{\partial F_1}{\partial x} - \frac{\partial F_2}{\partial y} \right] dA \quad F_1 = \frac{1}{2} x \quad F_2 = \frac{1}{2} y$

Masse - Masse-senter - Trehetsmoment

	Kurve	Flate	Volum
Masse	$M = \int_C \delta ds$	$M = \iint_S \delta dS$	$M = \iiint_V \delta dV$
Masse-senter	$\bar{x} = \frac{1}{M} \int_C x \delta ds$ $\bar{y} = \frac{1}{M} \int_C y \delta ds$ $\bar{z} = \frac{1}{M} \int_C z \delta ds$	$\bar{x} = \frac{1}{M} \iint_S x \delta dS$ $\bar{y} = \frac{1}{M} \iint_S y \delta dS$	$\bar{x} = \frac{1}{M} \iiint_V x \delta dV$ $\bar{y} = \frac{1}{M} \iiint_V y \delta dV$ $\bar{z} = \frac{1}{M} \iiint_V z \delta dV$
Trehetsmoment	$I_x = \int_C (y^2 + z^2) \delta ds$ $I_y = \int_C (x^2 + z^2) \delta ds$ $I_z = \int_C (x^2 + y^2) \delta ds$ $I_L = \int_C r^2 \delta ds$ $R_L = \sqrt{\frac{I_L}{M}}$	$I_x = \iint_S y^2 \delta dS$ $I_y = \iint_S x^2 \delta dS$ $I_L = \iint_S r^2 \delta dS$ $I_o = \iint_S (x^2 + y^2) \delta dS$ $R_L = \sqrt{\frac{I_L}{M}}$	$I_x = \iiint_V (y^2 + z^2) \delta dV$ $I_y = \iiint_V (x^2 + z^2) \delta dV$ $I_z = \iiint_V (x^2 + y^2) \delta dV$ $I_L = \iiint_V r^2 \delta dV$ $R_L = \sqrt{\frac{I_L}{M}}$

Fourier

Fourier-rekke (Periode 2L)	$f(t) = \frac{a_0}{2} + \sum_{n=1}^{\infty} \left[a_n \cos \frac{n\pi t}{L} + b_n \sin \frac{n\pi t}{L} \right]$ $a_n = \frac{1}{L} \int_{-L}^L f(t) \cos \frac{n\pi t}{L} dt \quad n = 0, 1, 2, 3, \dots$ $b_n = \frac{1}{L} \int_{-L}^L f(t) \sin \frac{n\pi t}{L} dt \quad n = 1, 2, 3, \dots$	Alternativt $a_n = \frac{1}{L} \int_c^{c+2L} f(t) \cos \frac{n\pi t}{L} dt \quad n = 0, 1, 2, 3, \dots$ $b_n = \frac{1}{L} \int_c^{c+2L} f(t) \sin \frac{n\pi t}{L} dt \quad n = 1, 2, 3, \dots$
Even funksjon $f(-t) = f(t)$	$a_n = \frac{2}{L} \int_0^L f(t) \cos \frac{n\pi t}{L} dt \quad n = 0, 1, 2, 3, \dots$ $b_n = 0 \quad n = 1, 2, 3, \dots$	
Odd funksjon $f(-t) = -f(t)$	$a_n = 0 \quad n = 0, 1, 2, 3, \dots$ $b_n = \frac{2}{L} \int_0^L f(t) \sin \frac{n\pi t}{L} dt \quad n = 1, 2, 3, \dots$	

Differensial-ligninger

Svinge-ligning	$m \frac{d^2 x}{dt^2} + c \frac{dx}{dt} + kx = F$	$mx_{tt} + cx_t + kx = F$
Bølge-ligning	$\frac{\partial^2 y}{\partial t^2} = v^2 \frac{\partial^2 y}{\partial x^2}$	$y_{tt} = v^2 y_{xx}$
Varme-ligning	$\frac{\partial u}{\partial t} = \kappa \nabla^2 u$	$u_t = \kappa y_{xx}$

Svinge-ligning m / initialbetingelser

<p>Kloss + fjær uten dempning</p>	$mx''+kx = F(t)$	$0 \leq t \leq L$ $x(0) = 0$ $x(L) = 0$
	$x(t) = x_0(t) + x_p(t) = c_1 \cos(\omega_0 t) + c_2 \sin(\omega_0 t) + x_p$ $F(t) = \sum_{n=1}^{\infty} b_n \sin \frac{n\pi t}{L}$ $b_n = \frac{2}{L} \int_0^L F(t) \sin \frac{n\pi t}{L} dt$ $x(t) = \sum_{n=1}^{\infty} B_n \sin \frac{n\pi t}{L}$ $= \sum_{n=1}^{\infty} \frac{1}{k - m \frac{n^2 \pi^2}{L^2}} b_n \sin \frac{n\pi t}{L}$	$\omega_0 = \sqrt{\frac{k}{m}}$
<p>Kloss + fjær med dempning</p>	$mx''+cx'+kx = F(t)$	$0 \leq t \leq L$ $x(0) = 0$ $x(L) = 0$
	$x_p(t) = \frac{F_0}{\sqrt{(k - m\omega^2)^2 + (c\omega)^2}} \sin(\omega t - \alpha)$ $\alpha = \tan^{-1} \frac{c\omega}{k - m\omega^2}$	$F(t) = F_0 \sin(\omega t)$
	$F(t) = \sum_{n=1}^{\infty} b_n \sin \frac{n\pi t}{L}$ $b_n = \frac{2}{L} \int_0^L F(t) \sin \frac{n\pi t}{L} dt$ $x(t) = \sum_{n=1}^{\infty} \frac{b_n \sin(\omega_n t - \alpha_n)}{(k - m\omega_n^2)^2 + (c\omega_n)^2}$	

Varme- og bølge-ligning m / initialbetingelser

<p>Varme-ligning m / initialbetingelser</p> <p>Fourier sinus-rekke</p>	$\frac{\partial u}{\partial t} = k \frac{\partial^2 u}{\partial x^2} \quad 0 < x < L$	$u(0,t) = 0$ $u(L,t) = 0$ $u(x,0) = f(x)$
	$u(x,t) = \sum_{n=1}^{\infty} b_n e^{-\frac{n^2 \pi^2 k}{L^2} t} \sin \frac{n \pi x}{L}$ $b_n = \frac{2}{L} \int_0^L f(x) \sin \frac{n \pi x}{L} dx$	
<p>Bølge-ligning m / initialbetingelser</p> <p>Fourier sinus-rekke</p>	$\frac{\partial^2 y}{\partial t^2} = a^2 \frac{\partial^2 y}{\partial x^2} \quad 0 < x < L$	$y(0,t) = 0$ $y(L,t) = 0$ $y(x,0) = f(x)$ $y_t(x,0) = g(x)$
	$y(x,t) = \sum_{n=1}^{\infty} A_n \cos \frac{n \pi a t}{L} \sin \frac{n \pi x}{L}$ $A_n = \frac{2}{L} \int_0^L f(x) \sin \frac{n \pi x}{L} dx$	<p>Problem A</p> $y(0,t) = 0$ $y(L,t) = 0$ $y(x,0) = f(x)$ $y_t(x,0) = 0$
	$y(x,t) = \sum_{n=1}^{\infty} B_n \sin \frac{n \pi a t}{L} \sin \frac{n \pi x}{L}$ $B_n = \frac{2}{n \pi a} \int_0^L g(x) \sin \frac{n \pi x}{L} dx$	<p>Problem B</p> $y(0,t) = 0$ $y(L,t) = 0$ $y(x,0) = 0$ $y_t(x,0) = g(x)$
	$y(x,t) = y_A(x,t) + y_B(x,t)$	